arXiv:1512.06310v1 [math.NT] 20 Dec 2015

A STERN-TYPE CONGRUENCE FOR THE SCHRÖDER NUMBERS

HUI-QIN CAO AND HAO PAN

Abstract. For the Schröder number

$$S_n=\sum_{k=0}^{n}\binom{n}{k}\binom{n+k}{k}\frac{1}{k+1},$$

we prove that

$$S_{n+2^\alpha}\equiv S_n+2^{\alpha+1}\pmod{2^{\alpha+2}},$$

where $n\geq 1$ and $\alpha\geq 1$.

In combinatorics, the Schröder number S_n is the number of all lattice paths from $(0,0)$ and (n,n) which just use the steps $(1,0)$, $(0,1)$, $(1,1)$ and contain no point above the line $y=x$. Using a simple combinatorial discussion, we can easily prove that

$$S_n=\sum_{k=0}^{n}\binom{n+k}{2k}\binom{2k}{k}\frac{1}{k+1}.$$

In fact, assuming a Schröder path exactly uses $n-k$ steps $(1,1)$, there are $\binom{n+k}{n-k}$ ways to insert those $n-k$ steps $(1,1)$ between k steps $(1,0)$ and k steps $(0,1)$. Also, it is well-known that the number of the lattice paths from $(0,0)$ and (k,k), which use the steps $(1,0)$, $(0,1)$ and never rise above the line $y=x$, is the Catalan number $C_k=\frac{1}{k+1}\binom{2k}{k}$. Furthermore, we know that the generating function for the Schröder numbers is

$$\sum_{n=0}^{\infty}S_nx^n=\frac{1-x-\sqrt{1-6x+x^2}}{2x}.$$

Nowadays, the combinatorial properties of the Schröder numbers have been widely discussed. For example, in [1, Exercise 6.39], Stanley listed many different combinatorial interpretations for S_n.

However, seemingly the arithmetical properties of S_n were seldom studied before. In fact, we know that S_n is always even for $n\geq 1$. The number

$$s_n=\frac{1}{2}S_n$$

2010 *Mathematics Subject Classification.* Primary 11B65; Secondary 05A10, 11A07.

Key words and phrases. Congruence; Schröder number.

is also called the little Schröder number, which counts the number of ways of dissecting a convex polygon with $n+1$ sides into smaller polygons by inserting diagonals. And in [2], Sun obtained several interesting congruences on the sum $\sum_{k=0}^{p-1} S_k m^{-k}$ modulo odd prime p.

On the other hand, the Euler number E_n is given by

$$\sum_{n=0}^{\infty} \frac{E_n}{n!} x^n = \frac{2}{e^x + e^{-x}}.$$

Clearly $E_n = 0$ if n is odd. The Euler numbers play an important role in number theory, since $E_{2n}/2$ coincides with the Dirichlet L-function $L(-2n, \chi_{-4})$. As early as 1851, Kummer had proved

$$E_{2n+(p-1)} \equiv E_{2n} \pmod{p} \tag{1}$$

for any odd prime p. Furthermore, in combinatorics, $(-1)^n E_{2n}$ is the number of all alternative permutations on $\{1, 2, \ldots, 2n\}$, i.e.,

$$(-1)^n E_{2n} = \#\{\sigma : \sigma(1) < \sigma(2) > \sigma(3) < \cdots > \sigma(2n-1) < \sigma(2n)\}.$$

In 1875, Stern proved another curious identity for the Euler numbers:

$$E_{2n+2^\alpha} \equiv E_{2n} + 2^\alpha \pmod{2^{\alpha+1}}, \tag{2}$$

where $\alpha \geq 1$. Clearly Stern's congruence implies that $E_{2n} \equiv E_{2m} \pmod{2^\alpha}$ if and only if 2^α divides $2n - 2m$. For the history and a proof of Stern's congruence, the reader may refer to Wagstaff's expository article [3].

Obviously from the viewpoint of either combinatorics or number theory, the Schröder numbers have no connection with the Euler numbers. However, in this short note, we shall show that S_n also satisfies a very similar congruence as (2).

Theorem 1. *For any $n \geq 1$ and $\alpha \geq 1$,*

$$S_{n+2^\alpha} \equiv S_n + 2^{\alpha+1} \pmod{2^{\alpha+2}}. \tag{3}$$

Clearly (3) is equivalent to

$$s_{n+2^\alpha} \equiv s_n + 2^\alpha \pmod{2^{\alpha+1}}, \tag{4}$$

i.e., $s_n \equiv s_m \pmod{2^\alpha}$ if and only if 2^α divides $n - m$. In fact, we can prove a stronger result.

Theorem 2. *For any $n \geq 1$ and $\alpha \geq 2$,*

$$S_{n+2^\alpha} \equiv S_n + (-1)^{\lfloor \frac{n-1}{2} \rfloor} 2^{\alpha+1} \pmod{2^{\alpha+3}}, \tag{5}$$

where $\lfloor x \rfloor = \max\{n \in \mathbb{Z} : n \leq x\}$. And when $\alpha = 1$, we also have

$$S_{n+2} \equiv S_n + 4 \pmod{16}. \tag{6}$$

Proof. We need another explicit form of S_n:

$$S_n = \sum_{k=1}^{n} \frac{1}{n}\binom{n}{k}\binom{n}{k-1}2^k. \tag{7}$$

Of course, (7) is known. However, for the sake of the completeness, here we give a proof:

$$\begin{aligned}
&\sum_{k=1}^{n} \frac{1}{n}\binom{n}{k}\binom{n}{k-1}2^k = \sum_{k=1}^{n} \frac{1}{n}\binom{n}{k}\binom{n}{k-1}\sum_{j=0}^{k}\binom{k}{j}\\
=&\sum_{j=0}^{n} \frac{1}{n}\binom{n}{j}\sum_{k=j}^{n}\binom{n-j}{k-j}\binom{n}{k-1} = \sum_{j=0}^{n} \frac{1}{n}\binom{n}{j}\binom{2n-j}{n-1}\\
=&\sum_{k=0}^{n} \frac{1}{n}\binom{n}{k}\binom{n+k}{n-1} = \sum_{k=0}^{n} \frac{1}{k+1}\binom{n+k}{2k}\binom{2k}{k},
\end{aligned}$$

where the Chu-Vandemonde identity is used in the third step.

Denote

$$T(n,k) = \frac{1}{n}\binom{n}{k}\binom{n}{k-1}2^k.$$

In fact, $N(n,k) = T(n,k)/2^k$ is also called the Narayana number, which counts the number of the Calatan paths with k peaks from $(0,0)$ to (n,n). Then

$$S_n = \sum_{k=1}^{n} T(n,k).$$

It is clear that

$$T(n,k) \equiv 0 \pmod{2^k},$$

since the Narayana number

$$\frac{1}{n}\binom{n}{k}\binom{n}{k-1} = \binom{n+1}{k}\binom{n-1}{k-1} - \binom{n}{k}\binom{n}{k-1}$$

is an integer.

For $\alpha \geq 1$, clearly

$$T(n+2^\alpha,1) = \frac{1}{n+2^\alpha}\binom{n+2^\alpha}{1}\binom{n+2^\alpha}{0}\cdot 2 = \frac{1}{n}\binom{n}{1}\binom{n}{0}\cdot 2 = T(n,1), \tag{8}$$

and

$$\begin{aligned}
T(n+2^\alpha,2) =&\frac{1}{n+2^\alpha}\binom{n+2^\alpha}{2}\binom{n+2^\alpha}{1}\cdot 2^2\\
=&2(n(n-1)+2^{\alpha+1}n-2^\alpha+2^{2\alpha})\\
=&T(n,2)+2^{\alpha+2}n-2^{\alpha+1}+2^{2\alpha+1}.
\end{aligned} \tag{9}$$

Moreover, noting that

$$\binom{n+2^\alpha}{2}=\binom{n-1}{2}+\binom{n-1}{1}\binom{2^\alpha}{1}+\binom{2^\alpha}{2}\equiv\binom{n-1}{2}+2^{\alpha-1}\pmod{2^\alpha},\tag{10}$$

we get

$$\begin{aligned}T(n+2^\alpha,3)=&\frac{1}{3}\binom{n+2^\alpha-1}{2}\binom{n+2^\alpha}{2}\cdot 2^3\\ \equiv&\frac{2^3}{3}\left(\binom{n-1}{2}+2^{\alpha-1}\right)\left(\binom{n}{2}+2^{\alpha-1}\right)\\ =&\frac{2^3}{3}\left(\binom{n-1}{2}\binom{n}{2}+(n-1)^2 2^{\alpha-1}+2^{2\alpha-2}\right)\\ \equiv&T(n,3)+(n-1)2^{\alpha+2}+2^{2\alpha+1}\pmod{2^{\alpha+3}}.\end{aligned}\tag{11}$$

Thus, when $\alpha=1$, we have

$$\begin{aligned}S_{n+2}=&\sum_{k=1}^{n+2}T(n+2,k)\equiv\sum_{k=1}^{3}T(n+2,k)\pmod{2^4}\\ \equiv&T(n,1)+\big(T(n,2)+n2^3-2^2+2^3\big)+\big(T(n,3)+(n-1)2^3+2^3\big)\\ \equiv&\sum_{k=1}^{3}T(n,k)+(2n-1)2^3-2^2\equiv S_n+2^2\pmod{2^4}.\end{aligned}$$

Below assume that $\alpha\geq 2$. Since $2^\alpha\geq\alpha+2$, we get

$$T(n,k)\equiv T(n+2^\alpha,k)\equiv 0\pmod{2^{\alpha+3}}$$

for $k>2^\alpha$. So

$$S_{n+2^\alpha}=\sum_{k=1}^{n+2^\alpha}T(n+2^\alpha,k)\equiv\sum_{k=1}^{2^\alpha}T(n+2^\alpha,k)\pmod{2^{\alpha+3}}.$$

By the Chu-Vandemonde identity,

$$\binom{n+2^\alpha}{k}=\sum_{j=0}^{k}\binom{n}{k-j}\binom{2^\alpha}{j}.$$

Since

$$\binom{2^\alpha}{j}=\frac{2^\alpha}{j}\binom{2^\alpha-1}{j-1}\equiv 0\pmod{2^{\alpha-\nu_2(j)}},$$

where

$$\nu_2(j)=\max\{b\in\mathbb{N}:\,2^b\mid j\}.$$

Clearly $\nu_2(j) \leq \lfloor \log_2 k \rfloor$ for any $1 \leq j \leq k$. We obtain that

$$\binom{n+2^\alpha}{k} \equiv \binom{n}{k} \pmod{2^{\alpha-\lfloor \log_2 k \rfloor}}.$$

If k is odd, then

$$\begin{aligned} T(n+2^\alpha,k) =& \frac{1}{n+2^\alpha}\binom{n+2^\alpha}{k}\binom{n+2^\alpha}{k-1}\cdot 2^k \\ =& \frac{2^k}{k}\binom{n-1+2^\alpha}{k-1}\binom{n+2^\alpha}{k-1} \\ \equiv& \frac{2^k}{k}\binom{n-1}{k-1}\binom{n}{k-1} = T(n,k) \pmod{2^{\alpha+k-\lfloor \log_2 k \rfloor}}. \end{aligned}$$

And if k is even, then

$$\begin{aligned} T(n+2^\alpha,k) =& \frac{1}{n+2^\alpha}\binom{n+2^\alpha}{k}\binom{n+2^\alpha}{k-1}\cdot 2^k \\ =& \frac{2^k}{k-1}\binom{n+2^\alpha}{k}\binom{n-1+2^\alpha}{k-2} \\ \equiv& \frac{2^k}{k-1}\binom{n}{k}\binom{n-1}{k-2} = T(n,k) \pmod{2^{\alpha+k-\lfloor \log_2 k \rfloor}}. \end{aligned}$$

Thus

$$T(n+2^\alpha,k) \equiv T(n,k) \pmod{2^{\alpha+k-\lfloor \log_2 k \rfloor}}$$

for each $1 \leq k \leq 2^\alpha$. It is easy to verify $k - \lfloor \log_2 k \rfloor \geq 3$ when $k \geq 5$. Thus we have

$$\begin{aligned} S_{n+2^\alpha} \equiv& \sum_{k=1}^{4} T(n+2^\alpha,k) + \sum_{k=5}^{2^\alpha} T(n+2^\alpha,k) \\ \equiv& \sum_{k=1}^{4} T(n+2^\alpha,k) + \sum_{k=5}^{2^\alpha} T(n,k) \pmod{2^{\alpha+3}}. \end{aligned}$$

Clearly

$$\binom{2^\alpha}{j} = \frac{2^\alpha}{j}\binom{2^\alpha-1}{j-1} \equiv 0 \pmod{2^{\alpha-1}}$$

for $1 \leq j \leq 3$. And

$$\binom{2^\alpha}{4} = 2^{\alpha-2}\binom{2^\alpha-1}{3} \equiv 2^{\alpha-2} \pmod{2^{\alpha-1}}.$$

So

$$\binom{n+2^\alpha}{4} = \sum_{j=0}^{4}\binom{n}{4-j}\binom{2^\alpha}{j} \equiv \binom{n}{4} + 2^{\alpha-2} \pmod{2^{\alpha-1}}.$$

Also, by (10),

$$\binom{n-1+2^\alpha}{2} \equiv \binom{n-1}{2} \pmod{2^{\alpha-1}}.$$

It follows that

$$\begin{aligned} T(n+2^\alpha,4) =& \frac{1}{3}\binom{n+2^\alpha}{4}\binom{n-1+2^\alpha}{2}\cdot 2^4 \\ \equiv& \frac{2^4}{3}\left(\binom{n}{4}+2^{\alpha-2}\right)\binom{n-1}{2} \\ \equiv& T(n,4)+\binom{n-1}{2}2^{\alpha+2} \pmod{2^{\alpha+3}}. \end{aligned} \tag{12}$$

Combining (8), (9), (11) and (12), we get

$$\begin{aligned} \sum_{k=1}^{4} T(n+2^\alpha,k) \equiv& T(n,1)+\left(T(n,2)+n\cdot 2^{\alpha+2}-2^{\alpha+1}+2^{2\alpha+1}\right) \\ &+\left(T(n,3)+(n-1)2^{\alpha+2}+2^{2\alpha+1}\right)+\left(T(n,4)+\binom{n-1}{2}2^{\alpha+2}\right) \\ \equiv& \sum_{k=1}^{4} T(n,k)+2^{\alpha+1}(n^2+n-1) \\ \equiv& \sum_{k=1}^{4} T(n,k)+(-1)^{\lfloor\frac{n-1}{2}\rfloor}2^{\alpha+1} \pmod{2^{\alpha+3}}. \end{aligned}$$

Consequently,

$$\begin{aligned} S_{n+2^\alpha} \equiv& \sum_{k=1}^{4} T(n,k)+(-1)^{\lfloor\frac{n-1}{2}\rfloor}2^{\alpha+1}+\sum_{k=5}^{2^\alpha} T(n,k) \\ =& S_n+(-1)^{\lfloor\frac{n-1}{2}\rfloor}2^{\alpha+1} \pmod{2^{\alpha+3}}. \end{aligned}$$

□

References

[1] R. P. Stanley, *Enumerative combinatorics, Vol. 2*, Cambridge University Press, Cambridge, 1999.

[2] Z.-W. Sun, *On Delannoy numbers and Schröder numbers*, J. Number Theory, **131**(2011), 2387–2397.

[3] S. S. Jr. Wagstaff, *Prime divisors of the Bernoulli and Euler numbers*, Number theory for the millennium III, 357–374, A K Peters, Natick, MA, 2002.

Department of Applied Mathematics, Nanjing Audit University, Nanjing 210029, People's Public of China

E-mail address: caohq@nau.edu.cn

Department of Mathematics, Nanjing University, Nanjing 210093, People's Republic of China

E-mail address: haopan79@zoho.com